

# Real-Time Measurement of Outdoor Tobacco Smoke Particles

Neil E. Klepeis, Wayne R. Ott, and Paul Switzer  
*Stanford University, Stanford, CA*

## ABSTRACT

The current lack of empirical data on outdoor tobacco smoke (OTS) levels impedes OTS exposure and risk assessments. We sought to measure peak and time-averaged OTS concentrations in common outdoor settings near smokers and to explore the determinants of time-varying OTS levels, including the effects of source proximity and wind. Using five types of real-time airborne particle monitoring devices, we obtained more than 8000 min worth of continuous monitoring data, during which there were measurable OTS levels. Measurement intervals ranged from 2 sec to 1 min for the different instruments. We monitored OTS levels during 15 on-site visits to 10 outdoor public places where active cigar and cigarette smokers were present, including parks, sidewalk cafés, and restaurant and pub patios. For three of the visits and during 4 additional days of monitoring outdoors and indoors at a private residence, we controlled smoking activity at precise distances from monitored positions. The overall average OTS respirable particle concentration for the surveys of public places during smoking was approximately  $30 \mu\text{g m}^{-3}$ . OTS exhibited sharp spikes in particle mass concentration during smoking that sometimes exceeded  $1000 \mu\text{g m}^{-3}$  at distances within 0.5 m of the source. Some average concentrations over the duration of a cigarette and within 0.5 m exceeded  $200 \mu\text{g m}^{-3}$ , with some average downwind levels exceeding  $500 \mu\text{g m}^{-3}$ . OTS levels in a constant upwind direction from an active cigarette source were nearly zero. OTS levels also approached zero at distances greater than approximately 2 m from a single cigarette. During periods of active smoking, peak and average OTS levels near smokers rivaled indoor tobacco smoke concentrations. However, OTS levels dropped almost instantly after smoking activity ceased.

## IMPLICATIONS

This article is the first peer-reviewed publication of systematic measurements of OTS concentrations. The main conclusion from these data, that OTS levels can be substantial under certain conditions, is vital to the development of outdoor tobacco control policy. Because adequate information on OTS levels and human exposures has previously been lacking, the estimation of health risks associated with OTS has been hindered, and public discourse concerning OTS has been impaired. The present study also has shown that continuous, portable airborne particle monitors are suitable in OTS investigations across a range of locations and environmental conditions.

Based on our results, it is possible for OTS to present a nuisance or hazard under certain conditions of wind and smoker proximity.

## INTRODUCTION

Secondhand tobacco smoke (SHS), also called environmental tobacco smoke (ETS) or passive smoke, is defined as diluted and dispersed air pollutant emissions generated from the consumption of tobacco products. Emissions may be exhaled by a smoker (mainstream) or by leaving the burning tip of a cigarette or cigar (sidestream). When occurring outdoors, SHS is called outdoor tobacco smoke (OTS).

Indoor SHS has an established connection to adverse health outcomes in adults and children, such as asthma, respiratory infection, and lung cancer.<sup>1</sup> More recent work has shown an association between SHS exposure and reduced cognitive ability in children,<sup>2</sup> increased respiratory disease in adults from work exposure and increased cancer for people exposed at home as children,<sup>3</sup> increased coronary heart disease in women exposed at home or work,<sup>4</sup> and a general increase in mortality for persons living with smokers.<sup>5</sup> The U.S. Surgeon General's Report titled "The Health Consequences of Involuntary Exposure to Tobacco Smoke" concludes that there is no level of exposure to SHS without some associated risk,<sup>6</sup> and the California Air Resources Board (CARB) recently designated SHS as a "toxic agent,"<sup>7</sup> a classification also given to pure compounds such as arsenic or benzene.<sup>8</sup>

The body of evidence demonstrates clear harm from SHS exposure and supports the pursuit of exposure reduction policies. In 1995, California Assembly Bill 13 was passed, which effectively banned smoking inside eating and drinking establishments throughout California. Other state- or country-wide initiatives that ban smoking inside bars and restaurants have also been enacted.<sup>9</sup>

Cities and counties have just started to institute bans on outdoor smoking, such as those for parks and beaches.<sup>10</sup> Bans may be supportable because of the drift of OTS inside buildings or from the littering of cigarette butts. Outdoor smoking bans may also serve to discourage smoking behavior in general, by making it more difficult for smokers to find a place to light up or by preventing children from associating smoking with enjoyable outdoor activities. However, the ongoing debate over the appropriateness of outdoor bans from an exposure standpoint suffers from a lack of air monitoring data. To date, no data have been published in the archival literature on the systematic measurement of human exposure to OTS.

**Table 1.** Characteristics of five real-time airborne particle monitors used in the present work.

Monitor Type	Abbreviation	Description	References
Piezobalance	PZB	The model 3511 (Kanomax, Inc.) and model 8510 (TSI, Inc.) PZBs measure RSP mass in units of $\mu\text{g m}^{-3}$ by passing an air stream through a 3.5- $\mu\text{m}$ size-selective particle impactor and onto a vibrating piezoelectric crystal. The frequency change in the crystal is converted to an average particle concentration with a resolution of $\sim 10 \mu\text{g m}^{-3}$ , which we record automatically in 1-min intervals using a custom-built logging system.	Ott et al. <sup>16</sup> and Sem et al. <sup>17</sup>
Nephelometer	NEPH	The model M903 integrating nephelometer (Radiance, Inc.) uses a flash lamp and optical filter to measure a light scattering coefficient ("extinction coefficient") for particles drawn into the instrument at intervals as small as 2 seconds. The instrument does not include a size-selective inlet.	Brauer et al. <sup>18</sup> and Radiance Research <sup>34</sup>
Laser photometer	SIDEPAK	The model AM510 SIDEPAK (TSI) is a 90° light scattering system using a 670-nm laser diode that is precalibrated by the manufacturer using Arizona road dust to measure aerosol mass in units of $\text{mg m}^{-3}$ . In the present work, we equipped the monitor with a 2.5- $\mu\text{m}$ impactor and used the internal logger to record levels at intervals as small as 10 seconds. Before each monitoring visit, the SIDEPAK inlet flow rate was adjusted to 1.7 $\text{l min}^{-1}$ using a Gilibrator primary flow calibrator	TSI <sup>35</sup>
Laser counter	GRIMM	The model 1.108 laser counter (GRIMM, Inc.) internally records counts of airborne particles every minute in 14 size ranges from 0.3 to 20+ $\mu\text{m}$ with a resolution of 1 particle count per liter. It measures light photons from a semiconductor laser that have been scattered at an angle of $\sim 90^\circ$	Grimm Technologies <sup>36</sup>
Photoelectric aerosol sensor	PAS	The model PAS 2000CE photoelectric aerosol sensor (EcoChem, Inc.) takes advantage of the physics of PAH photoemission on the surface of particles. It uses UV light to ionize PAH on particles $< 1 \mu\text{m}$ in diameter and measures the resulting electrical charges. The instrument is precalibrated to internally record the mass concentration of PAH in units of $\text{ng m}^{-3}$ at intervals as small as 30 seconds. Because it measures particle-bound PAH, the PAS instrument may respond differently to aerosols that have comparable total mass concentrations but vary in their surface PAH content.	Ott and Siegmann <sup>23</sup> and EcoChem Analytics <sup>37</sup>

To meet this need, we performed OTS monitoring surveys and controlled OTS experiments in public outdoor locations and a private residential patio using state-of-the-art, real-time particle sensing instruments. These instruments were anticipated to be useful for pinpointing and understanding transient elevations in OTS pollution. We expect that the results of our study will be helpful to those involved in tobacco-related policy development, as well as to risk assessors and environmental epidemiologists.

## EXPERIMENTAL WORK

Although there are many potentially toxic compounds in both the gaseous and particle phases of SHS,<sup>8,11</sup> for the present work we used airborne particle concentrations to characterize SHS levels. The use of particles to indicate the presence of SHS is common practice.<sup>12</sup> Airborne particles comprise a significant portion of the sidestream and mainstream mass emissions from burning cigarettes and other tobacco products, and indoor particle concentrations associated with SHS are substantial.<sup>13</sup> The size range of SHS particles is approximately 0.02–2  $\mu\text{m}$ ,<sup>14</sup> so that all of the SHS particles fall within the fine particulate matter (PM<sub>2.5</sub>) and respirable suspended particle (RSP; also called particulate matter with diameters  $< 3.5 \mu\text{m}$ ) size ranges. When inhaled, these particles can deposit in the human lung. Other benefits of using particles to characterize SHS are that particle concentrations can be measured using standard techniques, particles have a direct association with adverse health effects, and there are existing health standards for time-averaged particle concentrations.<sup>15</sup>

Because many types of portable continuous monitors for airborne particles are currently available, we decided for the present study to use a range of different instruments to characterize dynamic OTS levels in the field and under controlled conditions. The simultaneous use of multiple monitors of the same type and of different types allowed us to achieve a high level of confidence in measured OTS levels and to perform intensive evaluations and comparisons of the instruments.

## Real-Time Monitors

We used 5 types of portable real-time airborne particle monitoring instruments to measure OTS concentrations at intervals ranging from 2 sec to 1 min. The monitor types included a piezoelectric microbalance (piezobalance [PZB]), a photoelectric aerosol sensor (PAS), and three light-scattering photometers: an integrating nephelometer (NEPH), a laser particle counter (GRIMM), and a laser diode photometer (SIDEPAK). A brief summary of the characteristics of each real-time particle monitoring instrument, along with references to the scientific literature or manufacturers' guides, is given in Table 1. We selected each instrument because of its sensitivity to tobacco smoke particles, rapid response time, portability, and/or proven reliability in the field. In addition to these instruments, we used a real-time hot wire anemometer to record airflow (0.01  $\text{m sec}^{-1}$  threshold), temperature, and relative humidity (RH) every minute (VelociCalc Model 8386, TSI).

**Table 2.** Native units and conversion factors for real-time particle monitoring instrument readings.

<sup>a</sup> Instrument	Native Units	<sup>b</sup> Conversion Factor from Native Units to RSP Mass Concentration ( $\mu\text{g m}^{-3}$ )			
		$\bar{x}$	CI <sub>90</sub>	s	s/ $\bar{x}$
PZB	$\mu\text{g m}^{-3}$	—	—	—	—
NEPH	$10^{-6} \text{ m}^{-1}$	$4.6 \text{ m}^2 \text{ g}^{-1}$	$\pm 0.4$	$0.78 \text{ m}^2 \text{ g}^{-1}$	0.17
SIDEPAK	$\text{mg m}^{-3}$	$3.3 \times 10^{-3} \text{ mg } \mu\text{g}^{-1}$	$\pm 0.3$	$0.53 \times 10^{-3} \text{ mg } \mu\text{g}^{-1}$	0.16
GRIMM	counts L <sup>-1</sup>	$6300 \text{ counts m}^3 (\mu\text{g L})^{-1}$	$\pm 800$	$160 \text{ counts m}^3 (\mu\text{g L})^{-1}$	0.25
PAS	$\text{ng m}^{-3}$	$0.83 \text{ ng } \mu\text{g}^{-1}$	$\pm 0.1$	$0.19 \text{ ng } \mu\text{g}^{-1}$	0.23

Notes: <sup>a</sup>Real-time airborne particle monitoring instrument abbreviations: PZB = Kanomax or TSI PZB; NEPH = Radiance integrating nephelometer; SIDEPAK = TSI Sidepak laser photometer; GRIMM = Grimm laser counter; PAS = Ecochem photoelectric aerosol sensor. <sup>b</sup>The sample mean ( $n = 12$ ) of conversion factors from native units to estimated RSP mass concentration units are given for readings of each realtime airborne particle monitoring instrument. Also given are the 90% confidence intervals for the sample mean.<sup>38</sup> Dividing the native units by the conversion factor gives RSP units of  $\mu\text{g m}^{-3}$ . The conversion factors were determined by comparing average particle measurements for fresh cigarette smoke emissions of the NEPH, SIDEPAK, GRIMM, and PAS instruments against those for the PZB instrument taken during 12 monitor collocation experiments where valid PZB readings were available. Abbreviations:  $\bar{x}$  = the sample mean; CI<sub>90</sub> = the 90% confidence interval for the sample mean; s = the sample standard deviation; s/ $\bar{x}$  = the relative standard deviation.

The PZB was designated as the reference particle mass monitor because it provides direct measurements of RSP mass concentrations and it has a long history of use with tobacco smoke. The PZB has been shown to agree well with reference pump- and filter-based RSP measurements. Ott et al.<sup>16</sup> provide a review of previous studies that evaluated the PZB, including one by Sem et al.,<sup>17</sup> who report PZB mass readings for tobacco smoke to be within 15% of filter-based samples. Based on 9 recent experiments that we performed in a 9-m<sup>3</sup> chamber using cigarettes and incense as sources, we found that average mass readings of an impactor-equipped PZB were within approximately 10% of average mass concentrations determined from cyclone mass filter samples ( $R^2 = 96\%$ ).

### Instrument Testing and Calibration

The NEPH, SIDEPAK, GRIMM, and PAS continuous monitors can be used to estimate RSP mass concentrations. However, it is essential to first calibrate them with respect to the specific aerosol under study. We tested, calibrated, and compared the monitoring instruments for a tobacco smoke source during a set of 14 side-by-side experiments in a 44-m<sup>3</sup> room of a residence. For each experiment, a single cigarette was lit and allowed to burn by itself (smolder smoked) for 4–10 min. Doors and windows were kept closed, except to clear smoke from the room in between experiments. The room SHS particle concentrations were measured during and after each cigarette burn period. We subtracted background particle levels, which were observed just before smoking began, from all of the readings. Over the range of relative humidities, which we measured during the experiments (40–70%), we found no influence of RH on measured particle levels.

We calculated one conversion factor for each of 12 experiments where valid PZB readings were available (see Table 2) by taking the ratio of the fresh 5-min average for the PAS, SIDEPAK, GRIMM, and NEPH readings to the fresh PZB 5-min average RSP mass levels measured during a period starting 5–10 min after smoking stopped (at which time concentrations were evenly mixed in the room). Background levels were subtracted before taking the ratio. Fresh levels were used to determine conversion factors, because OTS was expected to consist exclusively

of fresh emissions. Except for the GRIMM monitor, we used the raw readings of each instrument to determine the conversion factors. In the case of the GRIMM, we used the sum of all of the particles from the lowest measured diameter of 0.3  $\mu\text{m}$  up to 3  $\mu\text{m}$ , because tobacco smoke particles are expected to be in the 0.02–2  $\mu\text{m}$  range.<sup>14</sup> Although linear regressions between 1-min average PZB readings and the other instruments across all of the well-mixed concentrations (fresh and aged) showed generally good agreement on a per-experiment basis ( $R^2 = 80\text{--}99\%$ ), there was evidence of a nonlinear relationship in many cases. Our use of ratios of background-subtracted 5-min average particle concentrations to calculate conversion factors, rather than linear regressions, resulted in lower relative variation for conversion factors, likely because it minimized biases because of deposition, coagulation, or evaporation of tobacco smoke particles occurring over time.

The average conversion factors from NEPH and SIDEPAK native units to RSP units observed in the present study ( $4.6 \text{ m}^2 \text{ g}^{-1}$  and  $3.3 \times 10^{-3} \text{ mg } \mu\text{g}^{-1}$ , respectively) are similar to those determined by other investigators. For example, Brauer et al.<sup>18</sup> found a value of  $4.7 \text{ m}^2 \text{ g}^{-1}$  for the NEPH conversion to mass for cigarettes, and both Travers<sup>19</sup> and Lee<sup>20</sup> found values of  $\sim 3 \times 10^{-3} \text{ mg } \mu\text{g}^{-1}$  for the SIDEPAK conversion. Lee performed 14 laboratory calibration tests of the SIDEPAK using gravimetric PM<sub>2.5</sub> filter samples and a smoking machine. Previous investigators have also found good agreement between personal nephelometers (e.g., the MIE personal DataRam) and reference gravimetric methods when calibrated for the target aerosols and adjusted properly for high RH.<sup>21</sup> Personal nephelometers, which have been used by U.S. Environmental Protection Agency (EPA) and others to characterize particle exposures,<sup>22</sup> operate on principles similar to the SIDEPAK and NEPH light-scattering photometers used in the current study. Our SIDEPAK conversion factor corresponds with an internal “custom calibration factor” of approximately 0.3 (dimensionless), which is calculated by multiplying our result by 1000 and taking the reciprocal.

Unlike the other particle instruments, the PAS is expected to exhibit variation in response to RSP based on the polycyclic aromatic hydrocarbon (PAH) content of

particle emissions, and it only responds to particles <1  $\mu\text{m}$  in diameter.<sup>23</sup> However, evidence suggests that the PAS-measured PAH in cigarette smoke consistently tracks RSP mass across a range of cigarette types and smoking styles. Ding et al.<sup>24</sup> report that mainstream smoke for U.S. cigarettes contains 1–1.6  $\mu\text{g}$  of PAH per cigarette. The average PAS-to-RSP conversion factor of 0.83  $\text{ng } \mu\text{g}^{-1}$ , which we observed in the present study for the PAS monitor's response to smolder-smoked Marlboro cigarette emissions, implies that 0.083% of the emitted particle mass consists of particulate PAH. Our value for the conversion factor is similar to a value of 0.8  $\text{ng } \mu\text{g}^{-1}$  observed by Repace<sup>25</sup> in a casino and values of 1 and 0.8–1.3  $\text{ng } \mu\text{g}^{-1}$  observed in two of our previous studies,<sup>26,27</sup> which used an older version of the PAS monitor (Model PAS 1000i, EcoChem, Inc.). We found that the older PAS 1000i monitor's response had to be reduced by a factor of 10 relative to the PAS 2000CE because of the fact that the 1000i uses a krypton bromine ultraviolet (UV) lamp, whereas the 2000CE uses a mercury vapor UV lamp.

Our use of particles measured by the GRIMM in the 0.3- to 3- $\mu\text{m}$  range avoided interference from nontobacco sources of ultrafine particles (<0.1  $\mu\text{m}$ ) and large dust particles (>3  $\mu\text{m}$ ). The empirical GRIMM conversion factor of 6300 counts  $\text{m}^3$  ( $\mu\text{g L}$ )<sup>-1</sup> agrees well with a theoretical mean value of  $\sim$ 6500 counts  $\text{m}^3$  ( $\mu\text{g L}$ )<sup>-1</sup> (relative standard deviation [RSD] of 0.15), which we calculated from the particle count data by assuming spherical particles, a uniform distribution of particle sizes in each size bin, a particle density of 1.1  $\text{g cm}^{-3}$ , and a lognormal particle size distribution with a mass median diameter of 0.2  $\mu\text{m}$  and a geometric standard deviation of 2.<sup>14</sup>

We estimated the error associated with readings of a given monitor by computing the ratio of 1-min values for matched instruments of the same type. We also estimated the error associated with conversion of native PAS, GRIMM, NEPH, and SIDEPAK readings to RSP mass units by computing the ratio of the estimated 1-min average RSP mass units for each monitor to the native RSP mass values measured by the PZB. The results of these calculations showed generally good consistency for intrainstrument and interinstrument comparisons, with the bulk of errors <10–20%.

### On-Site Monitoring Visits

To establish typical OTS levels, we conducted 15 on-site field visits to 10 public outdoor locations containing smokers, including restaurant and pub patios, cafés, airport sidewalks, and a public park (see Tables 3 and 4 and the location schematics in Figure 1). These visits were designed so that we could measure the average particle exposure attributable to emissions from real smokers that might occur during a meal at an outdoor establishment or while waiting on a sidewalk or in some other public area.

During each on-site visit, we made real-time measurements of airborne particles using the GRIMM and/or the PAS instrument or the SIDEPAK instrument. We used the PZB as a supplemental instrument during a single visit. We used the GRIMM, PAS, and SIDEPAK for the visits because they are more portable and unobtrusive than the PZB and NEPH monitors. For each visit, we measured OTS levels during periods with active smoking. To provide

background levels, we also measured during times when no tobacco sources were active.

For nine of the visits (S1–S9), we measured OTS particle levels using the PAS and/or GRIMM while sitting or standing on each patio or sidewalk and observing the activity of nearby cigarette and cigar smokers, but, because patrons engaged in uncontrolled smoking, we were not able to make precise measurements of the distance between smokers and the monitoring instruments. The monitors were generally positioned at breathing height (4–6 ft) or table height ( $\sim$ 3 ft). The inlets of the GRIMM and PAS monitors were placed within 12 in. of each other where possible. The time spent near active smokers ranged from 0.5 to 3.4 hr per visit.

For three on-site visits to outdoor patios (OC1–OC3), we smoked or smolder-smoked cigarettes or cigars near the monitoring positions for smoking periods of 0.1 and 0.5 hr. We used the GRIMM and/or PAS to measure OTS particles during these visits.

Finally, during three site visits to sidewalk patios (OP1–OP3), we measured OTS levels using the SIDEPAK at precise distances from active cigarettes, which were either smolder smoked or human smoked, for periods ranging from 0.6 to 1.7 hr. We also measured temperature, air speed, and RH continuously during these visits.

### Matched Monitor Experiments

To quantify the relationship between distance from the smoker and OTS concentration, that is, the proximity effect, and to make direct comparisons between OTS and indoor SHS levels, we performed controlled experiments on four days (E1–E4) at a private residence (Tables 3 and 4 and BP1 in Figure 1) using pairs of matched PAS, NEPH, and GRIMM instruments at different distances from burning cigarettes. We smolder smoked successive cigarettes both on the outdoor patio and inside the residence. For most experiments, we made continuous measurements of air speed, temperature, and RH.

The E1 experiments consisted of six outdoor patio experiments on a single day in which a cluster of single PAS, NEPH, and GRIMM monitors were surrounded by five burning cigarettes at distances of 2, 4, or 6 ft and heights of 3–4 ft for periods of 10 min per experiment. The cigarettes were positioned in concentric pentagonal arrangements so that cigarettes surrounded the monitors at equal distances for each experiment. This arrangement was expected to diminish the impact of wind direction on measured concentrations. In addition to the six cigarette experiments, we conducted two experiments in which a single cigar was smoked for 20–30 min at a distance of 4 ft from the monitor cluster. For all of the experiments, a second, identical cluster of particle monitors, which was intended to provide continuous background levels, was positioned  $\sim$ 28 ft (8.5 m) from the first cluster and around the corner of the house.

For experiments E2–E4, we built two mobile particle monitoring assemblies containing PAS, NEPH, and SIDEPAK instruments fastened to wheeled chairs. On each day, we created seven to nine periods of smolder-smoked cigarette activity lasting 30–50 min, using three to five individual cigarettes burned successively. The monitoring inlets and burning cigarettes were both at an approximate

**Table 3.** Summary of OTS on-site surveys and experiments.

<sup>a</sup> Name	<sup>b</sup> Location(s)	<sup>c</sup> Sources	<sup>d</sup> Duration (hr)	<sup>e</sup> Overall Average OTS RSP Concentration ( $\mu\text{g m}^{-3}$ )
<i>On-site surveys with uncontrolled human smokers</i>				
S1	PP1, PP2	H, CG, C	3.3	64 (PAS)
S2	PP2	H, CG, C	1.0	50 (PAS)
S3	PP1	H, CG, C	1.3	29 (PAS), 51 (GRIMM)
S4	RC	H, C	3.0	6 (PAS)
S5	RC	H, C	3.4	10 (PAS)
S6	PP3	H, CG, C	1.1	30 (PAS), 42 (GRIMM)
S7	PP3	H, C	1.4	26 (GRIMM)
S8	AP	H, C	0.6	31 (PAS), 30 (GRIMM)
S9	AP	H, C	0.5	56 (PAS), 15 (GRIMM)
<i>On-site surveys with controlled smolder-smoked cigarettes or a controlled smoker</i>				
OC1	PP3	H, CG, C	0.5	62 (PAS) 17 (GRIMM)
OC2	PK	S, C	0.4	67 (PAS), 23 (GRIMM), 60 (PZB)
OC3	BP2	H, C	0.1	27 (GRIMM)
<i>On-site proximity experiments with controlled smolder-smoked cigarettes or a controlled smoker</i>				
OP1	SC1, SC3	S, C	1.7	133 (SIDEPAK)
OP2	RP	S, C	0.6	106 (SIDEPAK)
OP3	SC2	H, C	1.4	109 (SIDEPAK)
<i>Private patio experiments with controlled smolder-smoked cigarettes or machine-smoked cigars</i>				
E1	BP1	S, C	2.0	48 (PAS), 19 (GRIMM), 10 (NEPH)
E2	BP1	S, C	3.7	47 (PAS), 28 (GRIMM), 10 (NEPH)
E3	BP1	S, C	3.9	61 (PAS), 29 (GRIMM), 22 (NEPH)
E4	BP1	S, C	2.5	38 (PAS), 18 (GRIMM), 16 (NEPH)

*Notes:* <sup>a</sup>S1–S9 = on-site visits (surveys) to patios and sidewalk areas with human smokers; OC1–OC3 = on-site controlled visits (surveys) for which the investigators controlled the smoking or smolder smoking of one or more cigarettes or cigars near the monitors; OP1–OP3 = on-site proximity experiments with controlled smolder- or human-smoked cigarettes positioned at precise distances from the monitoring positions; E1–E4 = controlled experiments performed at a private residence (patio, living room, bedroom) with smolder-smoked cigarettes positioned at precise distances from two separate monitoring positions. <sup>b</sup>Codes refer to one of the outdoor locations listed in Table 4. SC1, SC2, and SC3 = sidewalk cafés; PP1, PP2, and PP3 = pub patios; RC = resort café; RP = restaurant patio; PK = park plaza; AP = airport sidewalk; BP1 and BP2 = private backyard patio. <sup>c</sup>H = human smoked, S = smolder smoked, CG = cigars, C = cigarettes or cigarillos. <sup>d</sup>Duration of the monitoring period during which OTS sources were intermittently or continuously active. <sup>e</sup>The estimated average OTS RSP concentration in  $\mu\text{g m}^{-3}$  determined by converted measurements of a PAS, GRIMM, NEPH or SIDEPAK instrument (indicated in parentheses) taken during times when cigarettes or cigars were active. Background levels were subtracted. PAS = Ecochem photoelectric aerosol sensor; GRIMM = Grimm laser particle counter; PZB = Kanomax or TSI PZB; NEPH = Radiance integrating nephelometer; SIDEPAK = TSI Sidepak laser photometer. Results for S1–S9 include time when smokers were intermittently active at a location. Results for OC1–OC3, OP1–OP3, and E1–E4 include times when a cigarette or cigar was smoked or smolder smoked by the investigators near the monitoring position. Although experiments E2–E4 included indoor SHS measurements, they were not included in the calculated average OTS particle concentrations shown in the table.

height of 3–4 ft. To provide accurate background levels, we measured particle concentrations during intermediate time periods with no cigarette activity, which were of similar duration as the smoking periods. For each period of smoking activity, the two monitoring assemblies were placed on opposite sides of the source at distances of 0.25, 0.5, 1, 2, or 4 m. On day 4, the PZB instrument was added to the suite of monitoring instruments.

Immediately after five to six periods of controlled outdoor cigarette combustion on the backyard patio (BP1 location; E2–E4 experiments), we moved the monitoring assemblies indoors and performed several experiments in the bedroom or living room of the residence. The design of the indoor experiments was nearly identical to the outdoor experiments, except that only distances of 0.25 and 0.5 m from the burning cigarette were monitored, and the experiments were performed inside the house where all of the exterior doors and windows were closed

during periods of smoking activity. In addition, for one of the two living room experiments, a small fan was introduced to explore the effect of controlled air directionality. The fan blew air at a rate of  $\sim 0.4 \text{ m sec}^{-1}$  from the source toward one set of monitors. The airflow because of the fan was approximately equal to the average ground-level outdoor airflow rate that we observed during the patio experiments and on-site surveys (see below).

## RESULTS AND DISCUSSION

Measured concentrations of OTS consistently showed sharp spikes in airborne particle levels during periods when cigars or cigarettes were active. The structure of the peaks could be observed using the NEPH and SIDEPAK instruments, which provided readings at intervals of 2 and 10 sec, respectively (see Figure 2). Some peaks exceeded  $1000 \mu\text{g m}^{-3}$ . Transitory peaks of this nature, which are seen in close proximity to activity

**Table 4.** Characteristics of OTS monitoring locations.

Site	Abbreviation	<sup>a</sup> Width (m)	<sup>a</sup> Depth (m)	<sup>b</sup> Building Height (m)	<sup>c</sup> Distance to Building (m)	<sup>c</sup> Distance to Street (m)	<sup>d</sup> No. of Tables or Benches	<sup>d</sup> Seating Capacity
Sidewalk café 1	SC1	12	5	7	2	5	10	22
Sidewalk café 2	SC2	5	5	6	1	5	3	10
Sidewalk café 3	SC3	26	25	9	3	22	38	133
Pub patio 1	PP1	6	9	6	3	5	6	25
Pub patio 2	PP2	12	9	9	5	15	20	50
Pub patio 3	PP3	15	12	3	5	12	15	100
Restaurant patio	RP	12	5	7	3.7	35	9	43
Resort café	RC	9	7	4	2	320	2	8
Park plaza	PK	26	30	7	16	12	9	61
Airport sidewalk	AP	56	4	3		4	12	> 50
Backyard patio 1	BP1	11	5	6	2	11	–	–
Backyard patio 2	BP2	6	4	2.4	2	300	–	–

Notes: See Figure 1 for schematics of each location. <sup>a</sup>The approximate width and depth of the sidewalk or patio area intended for sitting or standing that is associated with the location. <sup>b</sup>The approximate height of the building facade immediately adjacent to the sidewalk or patio at each location. <sup>c</sup>The approximate distances from the monitoring position to the front of the nearest building and to the nearest roadway (farthest monitoring position of all those used). <sup>d</sup>The approximate number of tables or benches that were present at each location during the day of monitoring and the estimated number of seats (maximum occupancy).

sources, have been attributed to “microplumes” by previous investigators,<sup>28</sup> who observed them within 2 m of indoor point sources of pollution. Microplumes are defined as thin concentrated streams of smoke, or some other air pollutant, that follow complex trajectories during periods of release. When the microplumes impinge on a monitor inlet, the monitor momentarily registers a high peak in concentration. Over time and at further distance from the source indoors, the microplumes dissipate, and pollution becomes well mixed in an interior space, persisting long after the source has been extinguished. In contrast to persistent and mixed indoor levels, which exhibit smooth rises and decays in concentration, OTS consists entirely of periods characterized by microplumes. There is no period where OTS is well mixed, and OTS disappears almost instantly when tobacco sources are extinguished.

We analyzed the OTS data in terms of raw concentration readings, 1-min average concentrations, and averages on a per-visit, per-experiment, or overall basis. All of the results presented are for periods of continuous (experiments) or intermittent (surveys) active smoking. Before averaging and data analysis, we subtracted background levels for each day’s worth of data from each monitor. We created a consistent and integrated database by calculating 1-min averages for each monitor and by converting the native units of each monitor into units of RSP mass concentration ( $\mu\text{g m}^{-3}$ ) using the mean conversion factors in Table 2. The quantitative discussion of variation in OTS levels during each monitoring episode refers to either peak values over intervals as low as 2 or 10 sec or to 1-min average levels. During nearly all of the outdoor monitoring periods on patios and sidewalks where RH was measured, it was fairly low, averaging  $\sim 40\%$  with a range of 20–65%. Therefore, correction of OTS levels because of high RH was deemed unnecessary. Where measured, outdoor temperatures averaged 26 °C with a range of 10–38 °C and outdoor ground-level wind speeds ( $\sim 1$  m above

ground) averaged 0.41 m sec<sup>-1</sup> with a range of  $\sim 0$ –1.2 m sec<sup>-1</sup>.


### Typical OTS Levels

Tables 3 and 5 contain overall average OTS particle mass concentrations for periods of smoking during the outdoor on-site field visits and for breakdowns by various factors. Note that the results presented in Table 5 are not meant to imply direct comparisons of concurrent measurements for the different instruments, because not all of the monitors were used during a given visit.

As determined from PAS instrument measurements during the on-site visits with natural and controlled smoking (S1–S6, S8–S9, and OC1–OC2), average estimated RSP mass concentrations of OTS particles on a given day ranged from 6 to 67  $\mu\text{g m}^{-3}$  with an overall average of 33  $\mu\text{g m}^{-3}$ . The estimated GRIMM RSP levels for similar visits to outdoor patios (S3, S6–S9, and OC1–OC3) ranged from 17 to 51  $\mu\text{g m}^{-3}$  with an average of 34  $\mu\text{g m}^{-3}$ . The PZB levels from a single visit with controlled smoking near the monitor (OC2) averaged 60  $\mu\text{g m}^{-3}$  (0.4-hr averaging period).

In general, the variation in 1-min average OTS levels (Table 5) was very high, with overall RSDs of 1.7 for the PAS and GRIMM instruments. This variation results from the occurrence of sharp spikes in the OTS concentration time series because of swirling microplumes. Peaks in 1-min average OTS levels during site visits were observed to reach as high as 300–600  $\mu\text{g m}^{-3}$  as measured by the PAS and GRIMM instruments.

The estimated RSP mass concentrations determined from PAS measurements in the present work may have been influenced by nontobacco sources or differences in PAH emissions for different types of tobacco products or smoking styles relative to what we used during the calibration experiments. Ott and Siegmann<sup>23</sup> report very different PAH concentrations for different combustion sources. In the current study, we found that the PAS monitor was more sensitive to



**Figure 1.** Rough schematic diagrams of patios and sidewalks where OTS particle levels were monitored in proximity to smokers or burning cigarettes. All of the patios had at least an open roof, and many were open on three sides. Broken lines represent open boundaries, and solid lines indicate a surrounding wall or an adjacent building. Tables and benches are represented by circles or ovals, and rectangles indicate doorways to buildings or an opening in the wall or fence surrounding a patio. The approximate positions of active smokers and monitors during one or more visits are indicated by the letters “S” and “M,” respectively. See Table 4 for dimensions and other characteristics of each OTS monitoring location.


some non-OTS particles, such as diesel exhaust and soot from some types of candles, than the other instruments, because these emissions can be high in PAH. We minimized bias in the PAS measurements caused by other sources by including only levels for the PAS when no non-OTS sources or unexplained concentrations were observed.

Despite possible interference from other sources, the general validity of the PAS results (and their applicability to estimating OTS RSP) is supported by their generally


good agreement with the estimated RSP levels derived from the GRIMM instrument. Some of the differences that we observed between the two instruments may have resulted from microplume effects, in which localized peaks in particle concentration occurred near only one monitor’s inlet at a given instant.

To facilitate direct comparisons to PAS measurements performed in other studies, the estimated RSP values reported here can be converted back to the native units


### Smolder–Smoked Cigarettes at a Residence (BP1)



### Smoker at a Sidewalk Cafe (SC2)



### Smolder–Smoked Cigarettes at a Sidewalk Cafe (SC1)





**Table 5.** Observed OTS particle concentrations during patio and sidewalk café on-site visits in RSP mass units ( $\mu\text{g m}^{-3}$ ).

Factor	PAS <sup>a</sup>				GRIMM <sup>a</sup>				PZB <sup>b</sup>			
	<i>N</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$
Overall <sup>c</sup>	852	33	55	1.7	339	34	57	1.7	9	60	46	0.8
Cigarettes <sup>d</sup>	434	16	32	2.0	178	25	43	1.7	9	60	46	0.8
Cigars and cigarettes <sup>e</sup>	418	50	66	1.3	161	43	67	1.6	—	—	—	—
Closed area <sup>f</sup>	326	52	69	1.3	75	51	75	1.5	—	—	—	—
Open area <sup>g</sup>	526	21	40	1.9	264	29	49	1.7	9	60	46	0.8

*Notes:* This table contains grouped descriptive statistics calculated from 1-min average OTS particle measurements observed during nine onsite visits S1–S9, where natural smoking of cigarettes and cigars by smokers occurred (intermittent smoking), and three onsite visits OC1–OC3, during which one or more cigarettes or cigars were smolder smoked by the investigators near the monitor(s) (continuous smoking). The RSP mass units for the PAS and GRIMM were estimated using conversion factors from native PZB RSP values based on the results of controlled collocation experiments using all of the monitors and a cigarette source (see text). Background levels were subtracted from all of the instrument measurements. The monitors are abbreviated as follows: PAS = Ecochem photoelectric aerosol sensor; GRIMM = Grimm laser particle counter; and PZB = Kanomax or TSI PZB; The abbreviations for statistics are: *n* = sample size of 1-min average values;  $\bar{x}$  = RSP sample mean in  $\mu\text{g m}^{-3}$ ; *s* = RSP sample standard deviation in  $\mu\text{g m}^{-3}$ ; and *s*/ $\bar{x}$  = RSP relative standard deviation (dimensionless). <sup>b</sup>The PAS and GRIMM were used together for 6 of 13 visits (see Table 3). <sup>c</sup>The PZB was only present at the OC2 visit (see Table 3). <sup>d</sup>Results taken over all 13 visits. <sup>e</sup>Results for time periods when only cigarettes were observed to be active. <sup>f</sup>Results for time periods when both cigars and cigarettes were observed to be active. <sup>g</sup>Closed areas are patios located at restaurants or pubs and enclosed with a fence or wall on all sides so that directional air flow was effectively impeded (PP1 and PP2). <sup>h</sup>Open areas were sidewalks, sidewalk cafés, or parks where, although there may have been trees, umbrellas, and low barriers, there was enough open space that a potential “street canyon” effect could occur whereby air flow was channeled across the patio because of the presence of surrounding buildings (SC1, SC2, SC3, BP2, PK, RC, RP, and AP).

(nanograms per meter cubed) of the PAS instrument by using the conversion factor of  $0.83 \text{ ng } \mu\text{g}^{-1}$  presented above. For example, the average per-visit particle-bound PAH concentrations measured during on-site surveys where smoking occurred were 5–56  $\text{ng m}^{-3}$  with an overall average of 27  $\text{ng m}^{-3}$ , which is similar to the particulate PAH concentrations reported by Ott and Siegmann<sup>23</sup> using the same PAS 2000CE monitor.

As shown in Table 5, overall average OTS concentrations for time periods when both cigarettes and cigars were active (50 and 43  $\mu\text{g m}^{-3}$  for PAS and GRIMM, respectively) were 40–70% higher than those when only cigarettes were active (16 and 25  $\mu\text{g m}^{-3}$ ). This result may have occurred because cigars are active over a longer period of time than are individual cigarettes. In addition, average OTS concentrations measured by the PAS and GRIMM instruments during visits to outdoor patios that were enclosed by fences or walls (PP1 and PP2 locations) were 50% and 43% higher, respectively, than those observed in more open areas (52 and 51  $\mu\text{g m}^{-3}$  vs. 21 and 29  $\mu\text{g m}^{-3}$ ). In the more open patios (SC1–SC3, BP2, PK, RC, RP, and AP locations in Figure 1), which may have

contained tables, chairs, umbrellas, and low fences, air could flow across the patio, perhaps influenced by a “street canyon” effect characterized by air movement in a consistent direction along building boundaries. In contrast, the enclosed patios had walls on four sides that protected patrons from wind and may have contained OTS emissions to a greater degree.

### Outdoor versus Indoor Concentrations

The 3 days of monitoring at a residence (E2–E4), during which parallel measurements were performed indoors and outdoors using the PAS, GRIMM, NEPH, and PZB instruments, provide data for direct comparisons between OTS levels and indoor SHS levels. Tables 3 and 6 summarize the average OTS and indoor SHS particle concentrations observed during periods of active smoking for these experiments. Figure 2A shows the complete time series of one set of experiments (E3) for the NEPH instrument.

The effect of accumulation of cigarette emissions indoors and the effect of room volume were plainly evident during the experiments. Although OTS concentrations

**Figure 2.** (A) Real-time OTS and indoor SHS RSP mass concentrations determined from raw 2-second NEPH instrument readings during a suite of patio experiments (E3) performed in the backyard of a residence using smolder-smoked cigarettes. Average RSP mass concentrations are shown for each period when cigarettes were active, indicated by solid horizontal bars, for both northerly and southerly monitoring positions at source-receptor distances of 0.25, 0.5, 1, and 2 m. The southerly average concentrations, shown in larger typeface, were consistently higher than the northerly ones for outdoor measurements, likely because the prevailing winds were in the southerly direction. Outdoor air speed averaged  $0.5 \text{ m sec}^{-1}$  on the patio during times that cigarettes were active. The indoor air speed was close to zero. (B) Real-time OTS RSP mass concentrations determined from raw 10-second SIDEPAK instrument readings during an on-site proximity experiment (OP3) performed on a sidewalk patio with a human smoker. Average mass concentrations during periods of smoking are indicated by solid horizontal bars. The distance of the monitor from the smoker, which ranged over four values between 0.5 and 3.7 m, is also given. Air speed averaged  $0.16 \text{ m sec}^{-1}$  during times that cigarettes were active. (C) Real-time OTS RSP mass concentrations determined from raw 10-second SIDEPAK instrument readings during an on-site proximity experiment (OP1) performed on a sidewalk patio where cigarettes were smolder smoked at five different distances from the instrument, ranging from 0.3 m (1 ft) to 2.7 m (9 ft). Average mass concentrations during periods of smoking are indicated by solid horizontal bars. During this set of experiments, wind was consistently blowing in a single direction along the sidewalk. All of the concentrations were monitored in the downwind direction, except for the second cigarette at 0.6 m, for which concentrations were monitored in the opposite (upwind) direction. Air speed averaged  $0.5 \text{ m sec}^{-1}$  during times that cigarettes were active.

**Table 6.** Observed OTS and indoor SHS particle concentrations during controlled experiments E1–E4 and OP1–OP3 in RSP mass units ( $\mu\text{g m}^{-3}$ ).

Factor	PAS <sup>a</sup>				GRIMM <sup>a</sup>				NEPH <sup>a</sup>				PZB <sup>a</sup>				SIDEPAK <sup>b</sup>			
	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$	<i>n</i>	$\bar{x}$	<i>s</i>	<i>s</i> / $\bar{x}$
Outdoor <sup>c</sup>	1029	50	113	2.3	1052	22	50	2.3	1052	15	26	1.7	–	–	–	–	220	120	181	1.5
Living room <sup>c</sup>	235	33	33	1.0	235	30	35	1.2	235	32	22	0.7	30	35	12	0.3	–	–	–	–
Bedroom <sup>c</sup>	22	46	48	1.0	22	106	105	1.0	22	95	80	0.8	11	105	74	0.7	–	–	–	–
[0.25, 0.5] m <sup>d</sup>	328	108	175	1.6	332	45	76	1.7	332	35	38	1.1	–	–	–	–	104	177	228	1.3
[0.5,1] m <sup>d</sup>	202	43	72	1.7	202	16	21	1.3	202	11	11	1.0	–	–	–	–	51	128	126	1.0
[1,2] m <sup>d</sup>	301	19	25	1.3	310	12	34	2.9	310	7	8	1.3	–	–	–	–	32	32	30	0.9
[2,4] m <sup>d</sup>	198	8	9	1.1	208	4	5	1.3	208	2	2	0.9	–	–	–	–	33	11	7	0.6
Northerly <sup>e</sup>	465	28	76	2.7	465	17	54	3.3	465	12	26	2.2	–	–	–	–	–	–	–	–
Southerly <sup>e</sup>	451	72	144	2.0	465	27	45	1.6	465	20	28	1.4	–	–	–	–	–	–	–	–
Downwind <sup>f</sup>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	52	175	238	1.4
Upwind <sup>f</sup>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	10	2.5	1.0	0.4

**Notes:** This table contains grouped descriptive statistics calculated from 1-min average OTS particle measurements observed during four controlled day-long experiments at a residence (E1–E4) and three on-site proximity experiments (OP1–OP3), for which distance from continuously active tobacco sources was recorded precisely. The RSP mass units for PAS, GRIMM, NEPH, and SIDEPAK instruments are estimated based on conversion factors to PZB RSP mass concentration units that were calculated from the results of controlled cigarette smoking experiments performed using the collocated monitoring instruments. Background levels were subtracted. The monitors are abbreviated as follows: PAS = Ecochem photoelectric aerosol sensor; GRIMM = Grimm laser particle counter; NEPH = Radiance integrating nephelometer; PZB = Kanomax PZB; and SIDEPAK = TSI laser photometer. The abbreviations for statistics are: *n* = sample size of 1-min average values;  $\bar{x}$  = RSP sample mean in  $\mu\text{g m}^{-3}$ ; *s* = RSP sample standard deviation in  $\mu\text{g m}^{-3}$ ; and *s*/ $\bar{x}$  = relative standard deviation (dimensionless). <sup>a</sup>The PAS, GRIMM, and NEPH were used together for the E1–E4 day-long experiments (see Table 3). The PZB was only used during the indoor portion of the E4 experiments. <sup>b</sup>The SIDEPAK was only used (by itself) during the OP1–OP3 proximity experiments (see Table 3). <sup>c</sup>The "Outdoor" row contains statistics calculated from OTS levels across all of the experiments. The "Living Room" and "Bedroom" rows contain indoor SHS results for the two indoor locations when the fan was off or monitors were upwind from the fan. Indoor SHS levels were only measured at distances of 0.25 and 0.5 m from the monitoring positions. Apart from the two rows labeled "Living Room" and "Bedroom," all of the rows in the table are for OTS levels only. <sup>d</sup>The distance from the source in four groupings for OTS levels only. [ or ] indicates left or right limit is inclusive, and ) indicates right limit is exclusive. <sup>e</sup>For three outdoor experiments on the residential patio (E2–E4), groups of monitors were placed in northerly and southerly directions. <sup>f</sup>For these outdoor results (OP1), the plume of the cigarette emissions was observed to move in a single direction for the entire duration of the experiment, either toward the monitor (Downwind results) or away from the monitor (Upwind results).

dropped immediately to background levels when the cigarette sources were extinguished, indoor SHS concentrations persisted at relatively high levels and slowly decayed for hours until the doors were opened to ventilate the house.

As expected, the smaller bedroom with a volume of 44 m<sup>3</sup> had larger average indoor SHS particle concentrations during smoking (105  $\mu\text{g m}^{-3}$  from PZB) than the living room (35  $\mu\text{g m}^{-3}$  from PZB), which had a volume of >400 m<sup>3</sup> (see Table 6). The average indoor SHS levels observed in this study were similar to those observed by Özkaynak et al.,<sup>29</sup> who report that secondhand smoke contributes approximately 30  $\mu\text{g m}^{-3}$  on average to indoor particle levels in homes. In the present study, we observed PZB particle mass peaks in the living room and bedroom of approximately 50 and 200  $\mu\text{g m}^{-3}$ , respectively, which are similar to peak values that we observed in previous real-time monitoring studies of cigar and cigarette smoking in homes.<sup>26,30</sup>

The average OTS particle concentrations that we observed during each experiment across all of the distances were 10–22  $\mu\text{g m}^{-3}$  for the NEPH, 18–29  $\mu\text{g m}^{-3}$  for the GRIMM, and 38–61  $\mu\text{g m}^{-3}$  for the PAS, with overall averages of 15, 22, and 50  $\mu\text{g m}^{-3}$ , respectively (as shown in Table 6). The overall average indoor SHS concentrations, when the fan was not operating and at distances of 0.25 and 0.5 m only, were 30–35  $\mu\text{g m}^{-3}$  in the living room and 46–106  $\mu\text{g m}^{-3}$  in the bedroom for the different types of instruments. The higher levels measured outdoors and lower levels measured indoors by the PAS instrument may be because of emission of different

numbers of fine particles containing PAH, resulting from different cigarette combustion conditions.

Although the overall average OTS particle levels were lower than the indoor SHS levels when taken over all of the distances, except for the PAS instrument, the OTS levels at distances <0.5 m were roughly equal to or greater than the average indoor living room levels for all of the instruments. In addition, during other experiments on sidewalk cafés or restaurant patios (OP1–OP3) where wind effects were evident, average OTS levels during smolder or human smoking for the SIDEPAK instrument were 106–133  $\mu\text{g m}^{-3}$  for all of the distances, which are close to the levels observed in the bedroom during smoking. For individual cigarettes smoked or smoldered at a sidewalk café within 0.5 m of the monitor (OP1 and OP3; see Figure 2, B and C), average OTS particle levels measured by the SIDEPAK instrument during smoking exceeded 200  $\mu\text{g m}^{-3}$  for several different cigarettes and 500  $\mu\text{g m}^{-3}$  for another cigarette, indicating that circumstances can sometimes lead to short-term OTS levels that substantially exceed typical indoor SHS levels.

### Wind Effect

The experiment in the living room of the residence, where a fan was used to blow the plume of a burning cigarette toward a set of monitors at an air speed of  $\sim 0.4 \text{ m sec}^{-1}$ , demonstrates how wind can elevate OTS levels in downwind directions (Figure 2A). For this particular experiment, the fan increased average NEPH levels during smoking by

approximately three times at a downwind monitor relative to an upwind monitor.

This effect is further illustrated by our observation that the two sets of monitors positioned on either side of the active cigarette sources on the outdoor residential patio recorded much different OTS particle levels. The average levels in the northerly direction were approximately 40–60% lower than those in the southerly direction (Table 6). From the time profiles for one set of measurements (Figure 2A), it is evident that outdoor levels could be higher than corresponding indoor (non-fan) levels in one direction but near zero in the opposite direction.

The clearest evidence that wind leads to extremely high OTS levels during smoking was provided by the results of the OP1 experiment at the first sidewalk café where six cigarettes were smolder smoked at five distances from the SIDEPAK monitor (Figure 2C). For this experiment, the wind was observed to consistently blow the smoke microplumes in a single direction at an average speed of  $0.5 \text{ m sec}^{-1}$  when cigarettes were active. Upwind levels were practically zero, whereas the average downwind particle levels during smoking were  $582 \mu\text{g m}^{-3}$  at 0.3 m, and even at 1.2–2.7 m they were still elevated above background by  $13\text{--}41 \mu\text{g m}^{-3}$ . The 10-sec spikes in the downwind OTS particle time series sometimes exceeded  $1500 \mu\text{g m}^{-3}$ .

### Proximity Effect


We observed a clear reduction in OTS levels as the distance from a tobacco source increased. Generally, average levels within 0.5 m from a single cigarette source were quite high and comparable to indoor levels, and OTS levels at distances greater than 1 or 2 m were much lower. However, during on-site proximity experiments OP1 and OP3, OTS was still detectable by the SIDEPAK at distances of approximately 3–4 m from a single cigarette on sidewalk patios. A NEPH instrument also registered slightly elevated particle concentrations at a distance of 8 m from a cluster of burning cigarettes and around the corner of the house during a backyard patio experiment (E1).

To summarize and quantify the proximity effect observed in our study, we fit curves to average OTS particle concentrations ( $y$ ) as a function of the distance from the source ( $x$ ). Figure 3 shows two curves with separate fits for data from the sidewalk cafés (OP1–OP3:  $y = 44.4 x^{-2} + 27 x^{-1} + 4.1$ ) and the backyard patio (E1–E4:  $y = -0.3 x^{-2} + 16.8 x^{-1} - 2.8$ ), where distances were measured precisely. Every point represents the overall average for a given distance across all of the smoking periods and instruments at a given type of location. The levels on the private patio were generally lower and dropped off by 1–2 m, whereas the café levels, where winds may have been stronger and/or more directional, started out approximately four times higher and did not entirely drop off by 4 m.

### Previous OTS Studies

Before the current study, few data on OTS levels have been available. In an unpublished study, the CARB measured 1- and 8-hr average nicotine concentrations, number of active cigarettes, and wind characteristics outside

## Proximity Effect



**Figure 3.** Overall average OTS RSP mass concentrations across all of the instruments as a function of proximity to the OTS source, calculated using levels measured during experiments on a backyard patio and two sidewalk cafés for which source proximity was recorded precisely (see E1–E4 and OP1–OP3 in Table 3). Background RSP levels were subtracted from all of the measurements. The backyard patio experiments used smolder-smoked cigarettes in an area shielded by fences and trees. The sidewalk café experiments used smolder- or human-smoked cigarettes.

an airport, a college, a government center, an office complex, and an amusement park.<sup>8</sup> Average OTS nicotine concentrations were strongly affected by counts of the number of smokers and moderately affected by the size of the smoking area and the measured wind speed. The observed 8-hr average OTS nicotine levels in locations with relatively stronger winds or a smaller number of smokers were  $\sim 0.1 \mu\text{g m}^{-3}$  or less. In locations with a larger number of smokers, the levels could reach  $1$  or  $3 \mu\text{g m}^{-3}$ . These OTS levels are in the middle range of observed indoor SHS nicotine levels, which can average from  $0.01$  to  $10 \mu\text{g m}^{-3}$ . Based on the CARB study, Californians who spend time close to outdoor smokers could potentially be exposed to OTS levels similar to those associated with indoor SHS concentrations.

The general findings of the CARB study are compatible with the findings of the current work. The CARB results establish the potential for relatively high OTS exposures in places where smokers congregate. The experiments in the current work go further to quantify potential exposures under specific wind and proximity conditions, focusing on single smokers. Extrapolation of our controlled experimental methods and results to multiple smokers is complex, because one must consider the relative positions between each source and the receptor. Generally, we would expect that exposure increases in proportion with the number of active smokers. The exact increase depends on the amount of time that the receptor spends downwind and at a given distance from each source.

### Incremental Contributions to 24-hr Total Exposure

It is useful to calculate per-cigarette 24-hr incremental exposure ( $IE_{24}$ ) concentrations for OTS, where  $IE_{24}$  is defined as the contribution of a given OTS-related event involving one cigarette to a person's 24-hr total particle exposure. For example, during on-site experiment OP1, we observed an average OTS particle concentration at a distance of 0.3 m from a cigarette of  $582 \mu\text{g m}^{-3}$  in the downwind direction. Because the cigarette lasted approximately 10 min, we calculate a per-cigarette 24-hr incremental exposure as follows:  $IE_{24} = 582 \mu\text{g m}^{-3} \times 10 \text{ min}/1440 \text{ min} = 4 \mu\text{g m}^{-3}$ . The calculation amounts to a weighting of the per-cigarette average concentration by the proportion of time that the cigarette lasts with respect to the 24-hr (1440 min) day.

The incremental exposure concept allows one to combine exposures for different events and to compare the total to health-related standards or other reference levels. For example, if a person experienced nine cigarette events over the course of their day (with each event similar to the one that occurred at 0.3 m in the OP1 experiments), then their overall 24-hr OTS particle exposure would be  $9 \times 4 \mu\text{g m}^{-3} = 36 \mu\text{g m}^{-3}$ . This exposure would just exceed the EPA 24-hr health-based ambient standard for fine particles, which is currently  $35 \mu\text{g m}^{-3}$ . Note that the EPA standard was devised for ambient air pollution, which is likely to have substantially different composition than tobacco smoke pollution. However, because secondhand smoke contains many toxic compounds, including carcinogens, it is likely that, at a given airborne particle concentration, OTS carries the greater risk.

### CONCLUSIONS

The measurement of OTS is a new area in terms of epidemiologic and human exposure investigations. The present work provides some of the first evidence that OTS levels can be substantial under certain conditions of wind and proximity. The major findings of our research are summarized below.

First, real-time particle instruments, especially those based on light scattering, are useful in characterizing the determinants of OTS levels, which fluctuate on a time scale of seconds. The different particle detection instruments provide consistent findings and support the general conclusion that significant OTS levels can occur near smokers.

Second, outdoor particle concentrations measured close to a cigar or cigarette exhibit multiple concentration spikes, or microplumes, which are similar to those that have been observed close to indoor particle sources.

Third, average OTS particle levels near active sources over the course of one or more cigarettes can be comparable with average well-mixed indoor SHS particle levels observed to occur in living rooms or bedrooms during smoking. Average OTS particle concentrations can reach hundreds of micrograms per meter cubed. Unlike indoor SHS levels, which decay slowly over a period of hours, OTS levels drop abruptly to zero when smoking ends.

Fourth, OTS levels are highly dependent on wind conditions. Upwind levels are likely to be very low,

whereas downwind OTS levels during periods of active smoking can be very large with 10-second peak levels at the closest positions potentially exceeding  $1500 \mu\text{g m}^{-3}$  and average levels over the duration of a single cigarette potentially exceeding  $500 \mu\text{g m}^{-3}$ .

Fifth, OTS levels are highly dependent on source proximity. Levels at 0.25–0.5 m can drop by half or more as the distance increases to 1–2 m. At distances  $>2$  m, levels near single cigarettes were generally close to background. The concentrations at different distances are influenced by wind conditions. We found that it was possible for there to be detectable OTS levels at downwind positions of  $\geq 4$  m from a single active cigarette. Also, as the number of active cigarettes increases, the distance at which OTS is detectable is likely to increase.

Sixth, in outdoor restaurant patios and parks, where there may be multiple smokers, between 8 and 20 cigarettes smoked sequentially could cause an incremental 24-hr particle exposure greater than a threshold level of  $35 \mu\text{g m}^{-3}$  for a person who is within 0.5 m of the smokers. This threshold level is the 24-hr EPA health-based standard for fine particles.

Our results demonstrate that OTS can be high during periods of smoking in locations where persons are near active smokers. Therefore, it is possible for OTS to present a nuisance or hazard under certain conditions. Examples of scenarios where OTS levels might be high include eating dinner with a smoker on an outdoor patio, sitting at a table next to a smoker at a sidewalk café, sitting next to a smoker on a park bench, or standing near a smoker outside a building. Children who accompany a smoking parent or guardian may experience substantial exposure. Outdoor restaurant or pub workers who spend a significant portion of their time within a few feet of active smokers are also likely to receive relatively large total OTS exposures over the course of a day, possibly exceeding the EPA 24-hr health standard for fine particles. If one is upwind from a smoker, levels most likely will be negligible. However, if the smoker's position changes or one spends time downwind from a smoker, then moving to a distance of  $>2$  m can reduce the likelihood of experiencing elevated particle exposure because of OTS. Future studies should measure OTS levels for dynamic situations with multiple smokers, including continuous measurements of personal OTS concentrations or biomarker levels for workers in outdoor locations.

Support for health-based OTS bans may lie in a potential acute effect on susceptible populations. Short-term OTS exposures might be life threatening for high-risk persons, because the human cardiovascular system is very sensitive to secondhand smoke.<sup>31</sup> A recent before-and-after smoking ban study showed a decreased chance of myocardial infarction when a ban was in place,<sup>32</sup> which suggests that there is an acute risk associated with SHS exposure for persons at increased risk of coronary heart disease or with known coronary artery disease.<sup>33</sup>

### ACKNOWLEDGMENTS

The research described in this article was funded by California Proposition 99 through contract number

3317SC between the University of California San Francisco and Stanford University. Support was also provided by an award to Stanford from the Flight Attendant Medical Research Institute. The authors appreciate the assistance of Dr. Lance A. Wallace, formerly of U.S. Environmental Protection Agency (EPA), who provided technical guidance in planning quality assurance tests and also assisted with data analysis. The authors thank EPA for providing valuable equipment support, James L. Repace for his thoughtful ideas and encouragement, and Michelle House for her help with planning the outdoor field surveys.

## REFERENCES

1. *Respiratory Health Effects of Passive Smoking: Lung Cancer and Other Disorders*; Technical Report EPA/600/6-90/006F; U.S. Environmental Protection Agency, Office of Research and Development: Washington, DC, 1992.
2. Yolton, K.; Dietrich, K.; Auinger, P.; Lanphear, B.P.; Hornung, R. Exposure to Environmental Tobacco Smoke and Cognitive Abilities among U.S. Children and Adolescents; *Environ. Health Perspect.* **2005**, *113*, 98-103.
3. Vineis, P.; Airolidi, L.; Veglia, F.; Olgiati, L.; Pastorelli, R.; Autrup, H.; Dunning, A.; Garte, S.; Gormally, E.; Hainaut, P.; Malaveille, C.; Matullo, G.; Peluso, M.; Overvad, K.; Tjonneland, A.; Clavel-Chapelon, F.; Boeing, H.; Krogh, V.; Palli, D.; Panico, S.; Tumino, R.; Bueno-Demesquita, B.; Peeters, P.; Berglund, G.; Hallmans, G.; Saracci, R.; Riboli, E. Environmental Tobacco Smoke and Risk of Respiratory Cancer and Chronic Obstructive Pulmonary Disease in Former Smokers and Never Smokers in the EPIC Prospective Study; *BMJ* **2005**, *330*, 265-266.
4. Kawachi, I.; Colditz, G.A.; Speizer, F.E.; Manson, J.E.; Stampfer, M. J. Willett, W.C.; Hennekens, C.H. A Prospective Study of Passive Smoking and Coronary Heart Disease; *Circulation* **1997**, *95*, 2374-2379.
5. Hill, S.; Blakely, T.; Kawachi, I. Mortality among "Never Smokers" Living with Smokers: Two Cohort Studies, 1981-4 and 1996-9; *BMJ* **2004**, *328*, 988-989.
6. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*; Technical Report; U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health: Washington, DC, 2006; available at <http://sfgate.com> (accessed 2006).
7. Kay, J. Secondhand Smoke Toxic Air Contaminant: State Air Resources Board Vote Paves Way for Possible New Limits on Tobacco Use; *San Francisco Chronicle*, January 27, 2006; available at <http://sfgate.com> (accessed 2006).
8. California Air Resources Board. *Proposed Identification of Environmental Tobacco Smoke as a Toxic Air Contaminant*; Technical Report; California Environmental Protection Agency, California Air Resources Board, Office of Environmental Health Hazard Assessment: Sacramento, CA, 2005; Chapter 5, pp. V6-V19.
9. Cowell, A. British Ban Indoor Smoking; *New York Times*, February 15, 2006; available at <http://nytimes.com> (accessed 2006).
10. Covarrubias, A. Calabasas Snuffs out Public Smoking; *LA Times* (online) January 21, 2006; available at <http://www.latimes.com/> (accessed 2006).
11. Jenkins, R.A.; Guerin, M.R.; Tomkins, B.A. *The Chemistry of Environmental Tobacco Smoke: Composition and Measurement*, 2nd ed.; Lewis Publishers: Boca Raton, FL, 2000.
12. Jaakkola, M.S.; Jaakkola, J.J. Assessment of Exposure to Environmental Tobacco Smoke; *Eur. Respir. J.* **1997**, *10*, 2384-2397.
13. Nazaroff, W.W.; Klepeis, N.E. Environmental Tobacco Smoke Particles. In Morawska, L.; Salthammer, T., Eds.; *Indoor Environment: Airborne Particles and Settled Dust*; Wiley-VCH: Weinheim, Germany, 2004; 245-274.
14. Klepeis, N.E.; Apte, M.G.; Gundel, L.A.; Sextro, R.G.; Nazaroff, W. W. Determining Size-Specific Emission Factors for Environmental Tobacco Smoke Particles; *Aerosol Sci. Technol.* **2003**, *37*, 780-790.
15. U.S. Environmental Protection Agency. *Air Quality Criteria for Particulate Matter*; Technical Report EPA 600/P-99/002aF-bF; U.S. Environmental Protection Agency: Washington, DC, 2004.
16. Ott, W.R.; Switzer, P.; Robinson, J. Particle Concentrations inside a Tavern before and after Prohibition of Smoking: Evaluating the Performance of an Indoor Air Quality Model; *J. Air & Waste Manage. Assoc.* **1996**, *46*, 1120-1134.
17. Sem, G.; Tsurubayashi, K.; Homma, K. Performance of the Piezoelectric Microbalance Respirable Aerosol Sensor; *Am. Indust. Hyg. Assoc. J.* **1977**, *38*, 580-588.
18. Brauer, M.; Hurtle, R.; Lang, B.; Ott, W. Assessment of Indoor Fine Aerosol Contributions from Environmental Tobacco Smoke and Cooking with a Portable Nephelometer; *J. Expos. Anal. Environ. Epidemiol.* **2000**, *10*, 136-144.
19. Travers, M. Roswell Park Cancer Institute, Buffalo, NY. Personal communication, 2005.
20. Lee, K. University of Kentucky, Lexington, KY. Personal communication, 2006.
21. Wu, C.; Delfino, R.; Floro, J.; Samimi, B.; Quintana, P.; Kleinman, M.; Liu, L. Evaluation and Quality Control of Personal Nephelometers in Indoor, Outdoor and Personal Environments; *J. Expos. Anal. Environ. Epidemiol.* **2005**, *15*, 99-110.
22. Williams, R.; Suggs, J.; Rea, A.; Leovic, K.; Vette, A.; Croghan, C.; Sheldon, L.; Rodes, C.; Thornburg, J.; Ejire, A.; Herbst, M.; Sanders, W. The Research Triangle Park Particulate Matter Panel Study: PM Mass Concentration Relationships; *Atmos. Environ.* **2003**, *37*, 5349-5363.
23. Ott, W.R.; Siegmann, H.C. Using Multiple Continuous Fine Particle Monitors to Characterize Tobacco, Incense, Candle, Cooking, Wood Burning, and Vehicular Sources in Indoor, Outdoor, and In-Transit Settings; *Atmos. Environ.* **2006**, *40*, 821-843.
24. Ding, Y.S.; Trommel, J.S.; Yan, X.J.; Ashley, D.; Watson, C.H. Determination of 14 Polycyclic Aromatic Hydrocarbons in Mainstream Smoke from Domestic Cigarettes; *Environ. Sci. Technol.* **2005**, *39*, 471-478.
25. Repace, J.R. Respirable Particles and Carcinogens in the Air of Delaware Hospitality Venues before and after a Smoking Ban; *J. Occup. Environ. Med.* **2004**, *46*, 887-905.
26. Klepeis, N.E.; Ott, W.R.; Repace, J.L. The Effect of Cigar Smoking on Indoor Levels of Carbon Monoxide and Particles; *J. Expos. Anal. Environ. Epidemiol.* **1999**, *9*, 622-635.
27. Ott, W.R.; Wilson, N.; Klepeis, N.E.; Switzer, P. Real-Time Monitoring of Polycyclic Aromatic Hydrocarbons and Respirable Suspended Particles from Environmental Tobacco Smoke in a Home. In *Proceedings of the International Symposium on Measurement of Toxic and Related Air Pollutants*, U.S. Environmental Protection Agency and A&WMA, VIP-39, NTIS PB94-209756, Durham, NC, May 3-6, 1994; pp 887-892.
28. McBride, S.J.; Ferro, A.R.; Ott, W.R.; Switzer, P.; Hildemann, L.M. Investigations of the Proximity Effect for Pollutants in the Indoor Environment; *J. Expos. Anal. Environ. Epidemiol.* **1999**, *9*, 602-621.
29. Özkaynak, H.; Xue, J.; Spengler, J.; Wallace, L.; Pellizzari, E.; Jenkins, P. Personal Exposure to Airborne Particles and Metals—Results from the Particle TEAM Study in Riverside, California; *J. Expos. Anal. Environ. Epidemiol.* **1996**, *6*, 57-78.
30. Ott, W.R.; Klepeis, N.E.; Switzer, P. Analytical Solutions to Compartmental Indoor Air Quality Models with Application to Environmental Tobacco Smoke Concentrations Measured in a House; *J. Air & Waste Manage. Assoc.* **2003**, *53*, 918-936.
31. Barnoya, J.; Glantz, S. Cardiovascular Effects of Secondhand Smoke: Nearly as Large as Smoking; *Circulation* **2005**, *111*, 2684-2698.
32. Sargent, R.P.; Shepard, R.M.; Glantz, S.A. Reduced Incidence of Admissions for Myocardial Infarction Associated with Public Smoking Ban: Before and after Study; *BMJ* **2004**, *328*, 977-980.
33. Pechacek, T.F.; Babb, S. How Acute and Reversible Are the Cardiovascular Risks of Secondhand Smoke; *BMJ* **2004**, *328*, 980-983.
34. Radiance Research. *Operation Procedures, M903 Nephelometer, ROM Version 2.37*; Radiance Research: Seattle, WA, 1999.
35. TSI. *Model AM510 SidePak Personal Aerosol Monitor User Guide, P/N 19804566, Rev. B*; TSI, Shoreview, MN, 2003.
36. Grimm Technologies, I. *1100 Series Operators Manual*; Grimm Technologies, Inc.: Douglasville, GA, 2000.
37. EcoChem Analytics. *User's Guide: Photoelectric Aerosol Sensor (PAS 2000CE) and Diffusion Charger (DC 2000CE), Edition 1.4*; EcoChem Analytics: League City, TX, 2005.
38. Hahn, G.J.; Shapiro, S.S. *Statistical Models in Engineering*. John Wiley & Sons: New York, NY, 1967; p. 74.

### About the Authors

Neil E. Klepeis is a consulting assistant professor in the Department of Civil and Environmental Engineering at Stanford University. Wayne R. Ott is a consulting professor in the Department of Civil and Environmental Engineering and a visiting scholar in the Department of Statistics at Stanford University. Paul Switzer is a professor in the Department of Statistics at Stanford University. Please address correspondence to: Neil E. Klepeis, 16475 Tarpey Road, Watsonville, CA 95076; phone: +1-831-768-9510; <http://klepeis.net>.

Copyright of *Journal of the Air & Waste Management Association* (1995) is the property of *Air & Waste Management Association* and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.